

SYSTEMATIC REVIEW ON GASTROINTESTINAL HELMINTHS OF DOMESTIC RUMINANTS IN ETHIOPIA

Tewodros FENTAHUN

Unit of Biomedical Sciences, College of Veterinary Medicine and Animal Sciences, University of Gondar, P.O.Box 196, Gondar, Ethiopia

✉ Email: tedyvet@gmail.com;  ORCID: 0000-0002-2955-5638

📄 Supporting Information

ABSTRACT: This systemic review was conducted to identify, critically assess, and bring together available data from primary researches conducted so far on gastrointestinal (GI) helminthes of domestic ruminants in Ethiopia. In the country, GI helminths of domestic ruminants have been identified; examined and informative statistics has been extracted since a few decades ago. For this review, relevant articles were retrieved from English databases: PubMed, Google Scholar, Science Direct, Web of Science and Scientific Information Database (SID). Additional studies were recognized by scanning the African Journal Online (AJOL) that includes the Ethiopian Veterinary Journal and Bulletin of Animal Health and Production. Out of retrieved (n=154) articles, thirty three (n=37) articles which fulfilled the eligibility criteria were selected. Accordingly, twenty three GI helminthes species which belong to the three classes of helminthes have been found to occur in domestic ruminants in the country. The main genera reported so far are *Haemonchus*, *Strongyloides*, *Trichostrongylus*, *Oesophagostomum*, *Bunostomum*, *Fasciola*, *Monezia* and *Paramphistomum* whereas, *Haemochus contortus*, *Monezia expansa* and *Fasciolahepatica* are the most frequently reported species from Nematode, Cestode and Trematode classes respectively. The overall GI helminths prevalence ranged from 2.3% to 100% were reported. Simple flotation, sedimentation, modified McMaster technique and faecal culture are the most common and routine diagnostic methods which have been used in the country. Management aspects like husbandry practices, climate and host influences are found to be the principal contributing factors that affect GI helminths infections. So far, the control of GI parasites in the country is mainly focusing on the use of anthelmintics. Consequently, due to the lack of effective control strategies, anthelmintics are exclusively used which result in anthelmintics resistance. Generally, occurrence, epidemiological features, realistic control strategies, common diagnostic procedures and frequently encountered species are reviewed. Finally, the relevance of epidemiological knowledge and the development of efficient, sustainable and conventional control measures which cover wider ago-climatic zones of the country are suggested for controlling GI helminths infections and should be assessed timely.

Keywords: Anthelmintics, Domestic ruminant, Ethiopia, Gastrointestinal helminthes.

INTRODUCTION

As a result of having different agro-ecological zones and favorable environmental situations in Ethiopia, the country is believed to be endowed numerous livestock species and suitable for livestock production. It has the largest livestock population in Africa (Tilahun and Schmidt, 2012; CSA, 2013). According to CSA (2013) report, an estimated statistics showed that about 54 million cattle, 25.5 million sheep and 24.06 million goats are found in the country. Of the total cattle population, 98.95% are local breeds and the remaining are hybrid and exotic breeds. Furthermore, 99.8% of the sheep and nearly all goat population of the country are local breeds (CSA, 2013).

However, diseases have numerous negative impacts on production and productivity. Among diverse animal diseases encountered in the country, helminthes infections remain one of the most important limiting factors and a bottlenecking production and productivity these days (Elsa et al., 2012). By chance, the gastrointestinal tract of ruminants harbors variety of parasites particularly helminthes which can cause both clinical and subclinical parasitism. As stated by Lebbie et al. (1994), GI helminthes infections are of a global concern for livestock industry, which have devastating impact in Sub-Saharan Africa in general and in Ethiopia in particular as a result of wider range of agro-ecological factors which are fitting for diversified hosts and parasite species. Hence, gastrointestinal (GI) helminthiasis has become among the most important diseases encountered by livestock sector of Ethiopia and has been considered to be one of the major constraints in the development of the sector (Regassa et al., 2006). In Ethiopia, helminthiasis is responsible for 25% mortality and 3.8% weight loss in highland sheep (Bekele et al., 1992).

REVIEW
 PII: S222877012000031-10
 Received: March 19, 2020
 Revised: August 31, 2020

According to Zahid et al. (2005), the helminthes infections of ruminants are mostly caused by nematodes like Strongyloides spp., Ostertagia Spp., Bunostomum and Trichuris spp.; Cestodes such as Moniezia spp., Taenia spp. and Trematodes such as Paramphistomum spp., Fasciola spp. and Shistosoma spp. Perry et al. (2002) reported that GI nematodes have been ranked highest on globalindex with Haemonchus contortus on top. Moreover, Trichostrongyloidea that include genera such as Haemonchus, Trichostrongylus, Cooperia, and Nematodirus, and the Strongyloidea and Ancylostomatoidea with Oesophagostomum and Bunostomum, are the economically most important and widely prevalent GI nematodes (Takele et al., 2013; Winter et al., 2018). In addition, such a pervasive occurrence of the metacestodes; Cysticercus ovis and Cysticercus tenuicollis; Hydatid cyst (E.granulosus); Fasciola hepatic and Fasciola gigantica are most prevalent in the country and considered to be of great economic importance (Lemma et al., 1985; Regassa et al., 2009; Nigatu et al., 2009; Kebede et al., 2009; Feyesa et al., 2010; Nigatu, 2010; Endale et al., 2013; Abebe et al., 2015; Beyene and Hiko, 2019).

There are many associated risk factors influencing the occurrence and epidemiology of GI helminthes including age, sex, weather condition and husbandry or management practices (Khan et al., 2009). Factors such as host age, physiological status, breed, parasite species involved, and the epidemiological patterns (husbandry practices and climate variables) determine the degree of infection (Tembely and Hansen, 1996; Menkir et al., 2007). In Ethiopia, several studies have been conducted on ruminant helminthiasis in various regions reporting a prevalence ranged from 2.3–100% (Fikru et al., 2006; Abebe and Esayas, 2001; Yirsaw and Zewdu, 2015).

A number of published research reports have been found on GI helminthes of domestic ruminants in Ethiopia. However, these reports are found in a separate and unorganized way. So, comprehensive and well organized documentation about GI helminthes of domestic ruminant in the country is essential to support researchers, professionals and policy-makers to develop further actions on the control and prevention strategies. Therefore, the aim of this systematic review was to identify, assess critically, and bring together available data from primary research conducted so far on gastrointestinal (GI) helminthes of domestic ruminants in Ethiopia.

MATERIALS and METHODS

Source, selection strategies and protocol

This systematic review was carried out on GI helminthes of Ethiopian ruminant using available electronic and non-electronic databases. The electronic search was used as the primary search method. The main electronic databases used were PubMed, Google Scholar, Science Direct, Web of Science, Scientific Information Database (SID) were accessed from University of Gondar, Ethiopia. Relevant studies have been identified from English databases in Pub Med, Google Scholar, Science Direct, Web of Science, Scientific Information Database (SID). Additional studies were recognized by scanning the African Journal Online (AJOL) that includes the Ethiopian Veterinary Journal and Bulletin of Animal Health and Production. Moreover, complete and congress articles like original descriptive studies (designated as cross-sectional study) in sheep, goats and cattle were also considered. Epidemiological parameters such as prevalence of GI helminthes infection among sheep, goat and cattle, and main contributing factors like age, sex, and geographical sub-regions of Ethiopia were considered thoroughly. An intensive data searching was made to collect available information (Figure 3). The searching strategies used were combining the phrases close to GI helminths in large and small ruminants in Ethiopia as indicated in Figure 1. Searches were restricted to peer-reviewed articles published nationally or internationally in English language.




Figure 1 - A diagram showing a strategy for words combination to access relevant articles.




Figure 2- Map of Ethiopia showing study sites of GI helminthes of domestic ruminants.




Figure 3 - Flow diagram describing the study design process and literature search on GI helminthes in domestic ruminants.

Data type and collecting methods

Published papers were scanned by quick reading to select relevant articles. Important articles were defined and included in the current review as one that contained information on GI helminthes of cattle, sheep and goat in Ethiopia. References of all relevant articles were searched to identify articles that were missed by the electronic search. Any

identified article was subjected to the same inclusion process as a data type. Following that all relevant articles were reviewed, extracted and compiled in a searchable database (Microsoft Access software, ver. 2007). Extracted information included authors name, study sites, year of the study commenced and ended, year of publication, study design, species of animals, sampling procedure, number of animals selected, body condition, sex, altitude of the study site, laboratory procedure to detect the parasite, testing methodology and prevalence of GI helminthes reported including their associated risk factors.

Reports collected on GI helminths in Ethiopia

The initial electronic searches yielded a total of (n=154) studies. After scrutinizing these and eliminating duplicates, thirty three (n=37) were considered (Figure 4) including thesis containing relevant information concerning GI helminths of ruminants in Ethiopia. This review work documented that of the relevant articles about 8% (3) were not retrieved by any of the search engines, 19% (7) were retrieved by a single search engine, 27% (10) by two, 16% (6) by three and 30% (11) by all four databases. Providing a balanced and an impartial summary of the topic using these representative studies was at the core of this review. No restrictions other than identifying ovine, caprine and bovine GI helminthes in Ethiopia were imposed on the inclusion criteria. This minimized literature selection bias and provided an exhaustive list of GI helminths. However, for factors associated with domestic ruminant GI helminthes in Ethiopia, strict inclusion criteria were applied in order to only select the critical studies. In order for the current review to be critically appraised, repeatability was crucial. This was achieved by documenting every step of the review process.




Figure 4 - Frequency of articles relevant to the systematic review on GI helminths of domestic ruminants in Ethiopia.

Occurrence and spatial distribution

The present review work revealed that the occurrences of GI helminthes infections were reported in different parts of the country (Figure 2). The prevalence of GI helminthes is summarized in table 1. The majority of the prevalence studies were specific on some helminthes genera like *Strongyles*, *Fasciola*, *Shistosoma* and some families like GI nematodes, Trematodes and Cestodes. However, adequate sources were not accessed compared to the number of studies done on specific species and families of the GI helminthes of domestic ruminants. In addition, most studies are based on the coprological examinations and few studies were conducted on abattoir surveys in the country.

According to the result obtained from the review, the overall prevalence of GI helminthes is ranged from 13.2% in Awash River Basin (Afar region) to 81.5% in Debre Zeit/Bishoftu (Bersissa et al., 2011; Moti et al., 2013). Furthermore, the review underlined that the majority of the identified species are found from nematodes and the lowest from Cestode. On the other hand, *Haemonchus contortus*, *Moneizia expansa* and *Fasciola hepatica* are the predominant species from Nematode, Cestode and Trematode classes respectively. Furthermore, *Nematodirus* spp., *Toxocaravitulorum* and *Schistosoma* spp. are less prevalent (Table 2). In Ethiopia the majority of the studies were focusing only on small ruminants' GI helminthes than large ruminants. Therefore, the highest prevalence of the parasite was reported from sheep (91.2%) in Gode (Ogaden), Southern Ethiopia than cattle (21.9%) in Nekemete in Western part of Ethiopia (Table 1). On the other hand, most Abattoir survey reports have also shown a status of GI helminths of cattle than small ruminants.

Table 1 - List of selected studies on GI helminthes of domestic ruminants in Ethiopia

Study site(s)	Study animals and Sample size				Study period	ST	Diagnostic technique	Prevalence (%)				Sample	Author(year)
	Sheep	Goat	Cattle	Total				Sheep	Goat	Cattle	Total		
Adama	92	208	852	1152	Nov2007-Apr2008	CS	PME	29.3	6.7	46.8	-	Liver	Getawa et al. (2010)
Adami Tulu	-	499	-	499	Jul1997-Jun1998	CS	MM	-	45.5	-	45.5	F	Etana (2002)
Addis Ababa	560	-	-	560	Sep 1985 - Jun 1986	CS	PME	19.6	-	-	19.6	Liver	Bekele et al.(1988)
Ambo	90	30	-	120	Feb-May 2013	CS	DS,FL,SD	47.8	53.3	-	-	F	Temesgen and Walanso (2015)
Arba Minch	241+20	357+25	-	598 + 45	Jan2010-Aug 2011	CS	SD,FL,MM,PME	79.7; 85	63.6; 68	-	70.1;	F, GIT	Nejib et al. (2014)
Asella	-	-	384	384	Nov 2013-Apr 2014	CS	FL	-	-	49	49	F	Addisu and Berihu (2014)
Asella	408	-	-	408	Nov 2008-Apr 2009	CS	MMM	68.1	-	-	68.1	F	Diriba et al. (2013)
Awash	3,697	-	-	3,697	Jan-Dec 2005	CS	EAC	13.2	-	-	13.2	F	Ahmed et al.(2007)
Bahir Dar	340	-	420	-	May 2005 - Dec2006	-	-	10.6	-	34.05	-	Liver	Nigatu et al. (2009)
Bahir Dar	384	-	384	768	Nov2010-March 2011	CS	SD, PME	10.9; 16.9	-	31.5; 45.3	-	F, Liver	Ayalew and Endalkachew (2013)
Bahir Dar	-	-	384	4134	Nov2008-March2009	CS	SD, PME	-	-	39.95	39.95	F, Liver	Fikirtemariam et al. (2013)
Bahir Dar	-	-	384	384	Nov2013-April 2014	CS	SD	32.3	-	-	32.3	F	Yitayal et al. (2015)
Bedele	-	-	500	500	Oct 2011-March 2012	CS	SD, FL, MM	-	-	64.2	64.2	F	Moti et al. (2013)
Bishoftu (Debre Zeit)	1152	1536	-	2688	Dec2005 - Jun 2006	CS	PME	58.5	43.8	-	-	Liver	Jibat et al. (2008)
Bishoftu (Debre Zeit)	157	65	-	222	Nov2007-Apr2008	CS	FL,SD, CC	81.0	83.0	-	81.5	F	Bersissa et al. (2011)
Bishoftu (Debre Zeit)	-	-	326	326	Nov2011-Apr 2012	CS	FL, SD,	-	-	61	61	F	Cheru et al. (2014)
Debre Berhan	2500	-	-	2500	1987	CS	SD,FL	18	-	-	18	F, PME	Njau et al.(1990)
Dembi Dolo	255	245	257	757	2003-2004	CS	DS, SD,FL, MM	75.3	84.1	50.2	69.6	F	Fikru et al.(2006)
Dessie	510	420	-	930	Nov2011-Mar 2012	CS	PME	21.04	27.61	-	-	OM, PE, Liver,MS	Abebe et al.(2015)
Dire Dawa	425	420	-	845	Nov2011-April 2012	CS	PME	22.8	26.4	-	24.6	Liver, PE OM	Endale et al. (2013)
Durbete	202-	-	330	532	Oct 2014- Apr 2015	CS	SD	2.3	-	24.6	26.9	F	Yirsaw and Zewdu (2015)
Gondar	458	100	-	558	Nov-Jan 2008	CS	SD, FL	46.07	55	-	47.67	F	Shimelis et al. (2011)
Gondar, Finote Selam, Injibara	-	-	22,755	22,755	Sept 2002-2007	CS	PME	-	-	79.5	79.5	Liver	Nigatu (2010)
Haremaya, Harar, Dire Dawa, Jijiga	655	632	-	1,287	May 2003-Apr 2005	CS	PME	-	-	-	-	GIT	Menkir et al. (2007)
Hawassa	284	226	-	510	Nov2008-Feb 2009	CS	MMM	47.2	38.9	-	-	F	Abebe et al. (2010)
Hawassa	-	-	632	632	Dec2008 - Mar 2009	CS	PME	-	-	52.6	52.6	Liver	Feyesa et al. (2010)
Hawassa	180	132	-	312	Jan-Jun 2006	CS	PME	91.1	87.1	-	-	A	Thomas et al. (2007)
Jimma	-	-	210	210	Nov2008-Apr 2009	CS	FL,SD, MMM	-	-	77.6	77.6	F	Hailu et al. (2011)
Kelala	230	-	154	384	Sept1997-Apr1998	CS	FL, SD	53	-	38	-	F	Tesfaye (1998)
Kombolcha	-	-	400	400	2011	CS	PME	-	-	17	17	Liver	Fufa et al. (2012)
Mekelle	-	-	1023	1023	Nov2007 - Feb2008	CS	PME	-	-	7. 23	7. 23	Liver	Getachew and Ashwani (2013)
Mekelle, Adigrat, Axum, Humera, Maichew, Shire	-	-	5,194	5,194	2007/8	CS	PME	-	-	22.1	22.1	Liver	Kebede et al. (2009)
Metehara, Semera, Jigjiga	92	91	-	183	Nov1998-Apr 1999	CS	MMM, PME	97.03	100	-	-	F, GIT	Abebe and Esayas(2001)
Nekemete	-	-	384	384	Nov2011-March 2012	CS	PME	-	-	21.9	21.9	GIT	Alula et al. (2013)
Ogaden-Gode	114	82	-	196	Aug2003-March 2004	CS	PME	91.2	82.9	-	-	A	Kumsa and Wossene (2006)
Wolaita Soddo	-	-	415	415	Nov2007 - Apr2008	CS	PME	-	-	11.3	-	Liver	Regassa et al. (2009)
Wukro	384	-	-	384	Nov 2013-Apr 2014	CS	PME	40.9	-	-	40.9	A	Lidya and Berihun (2015)

-: No information available; ST=Study design; PME=Postmortem examination; CS=Cross sectional; SD=Sedimentation, FL=Flotation, MM=Mc Master; DS=Direct Smear; MMM=Modified Mc Master; F= Faeces; A=Abomasum; GIT=Gastrointestinal content; EAC= Ethyl-acetate Centrifugation; PE=peritoneum; OM=Omentum; MS=Mesentry; RT= Retrospective

Table 2 - GI helminthes species/genus composition; collected from each selected studies in Ethiopia.

Nematode	Species of parasite Identified (%)											Location	Altitude of study area	Author (year)
	Species of animal			Cestode	Species of animal			Trematode	Species of animal					
	Sheep	Goat	Cattle		Sheep	Goat	Cattle		Sheep	Goat	Cattle			
-	-	-	852	Hydatid cyst	40.0	55.6	37.1	-	-	-	-	Adama	1770	Getaw et al. (2010)
<i>H. contortus</i>	-	30.3	-	<i>M. expansa</i>	-	14.6	-	-	-	-	-	Adami Tulu	900	Etana (2002)
Strongyloides spp.	-	16.4	-											
<i>Trichostrongylus</i> spp.	-	13.8	-											
<i>Trichuris</i> spp.	-	2.2	-											
<i>Nematodirus</i> spp.	-	2.8	-											
<i>Chabertia ovina</i>	-	5.0	-											
<i>Bunostomum</i> spp.	-	5.4	-											
-	-	-	-	<i>T. hydatigena and Hydatid cyst</i>	18.6	-	-	-	-	-	-	Addis Ababa	2,326	Bekele et al. (1988)
Strongyles spp.	47.8	53.3	-	-	-	-	-	<i>Fasciola</i> spp.	4.4	6.7	-	Ambo	2185	Temesgen and Walanso (2015)
<i>H. contortus</i>	A*	A*	-	-	Arba Minch	1400	Nejib et al. (2016)							
<i>Trichostrongylus</i> spp.	A*	A*	-	-										
<i>Teladorsagia</i> spp.	A*	A*	-	-										
<i>T. colubriformis</i>	A*	A*	-	-										
<i>Haemonchus</i> spp.	A*	A*	-	-										
<i>Ostertagia</i> spp.	-	-	1.8	Assela	1500-2300	Addisu and Berihu (2014)								
<i>Oesophagostomum</i> sp.	-	-	1.3											
Strongloid spp.	-	-	2.6											
<i>Trychostrongylus</i> spp.	-	-	3.6											
<i>Hemonchus</i> spp.	-	-	11.7											
<i>Bunostomum</i> spp.	-	-	4.4											
Strongyles spp.	64.0	-	-	Asella	2500-3000	Diriba et al. (2013)								
Strongyloides spp.	7.4	-	-											
<i>Trichuris</i> spp.	3.7	-	-											
-	-	-	-	-	-	-	<i>Fasciola</i> spp.	13.2%	-	-	Awash	550-850	Ahmed et al. (2007)	
-	-	-	-	<i>E. granulosus</i>	25.0	-	16.1	-	-	-	Bahir Dar	1784	Nigatu et al. (2009)	
-	-	-	-	-	-	-	-	<i>F. hepatica and F. gigantica</i>	16.9	-	45.31	Bahir Dar	1784	Ayalew and Endalkachew (2013)
-	-	-	-	-	-	-	-	<i>F. hepatica</i>	-	-	89.7	Bahir Dar	1784	Fikirtemariam et al. (2013)
-	-	-	-	-	-	-	-	<i>F. gigantica</i>	-	-	3.63			
-	-	-	-	-	-	-	-	Mixed	-	-	6.67			
-	-	-	-	-	-	-	-	<i>F. hepatica</i>	-	-	32.3	Bahir Dar	1784	Yitayal et al. (2015)
<i>Trichuris</i> spp.	-	-	6.8	-	-	-	-	<i>Fasciola</i> spp	-	-	51.4	Bedele	2060	Moti et al. (2013)
<i>Trichostrongyles</i> spp.	-	-	11.0	-	-	-	-	<i>Paramphistomum</i> spp,	-	-	18.7			
-	-	-	-	<i>St. hepatica</i>	9.5	12.1	-	<i>Fasciola</i> spp.	6.9	3.6	-	Bishoftu	2155	Jibat et al. (2008)
-	-	-	-	<i>C. tenuicollis</i>	5.2	8.3	-	-	-	-	-	-	-	-
Strongyle type	56.60	61.0	-	<i>M. expansa</i>	13	29.2	-	<i>Fasciola</i> spp.	0.63	0	-	Bishoftu	2155	Bersissa et al. (2011)
Strongyloides spp.	8.2	15.4	-	-	-	-	-	-	-	-	-	-	-	-

Trichuris spp.	5.0	0.0	-											
Strongyle spp.	-	-	41.4											
Trichuris spp.	-	-	41.4											
Toxocara spp.	-	-	5.2	<i>Monezia</i> spp.	-	-	2.8	<i>Fasciola</i> spp.	-	-	36.5	Bishoftu	2155	Cheru et al. (2014)
Trichuris spp.	-	-	5.2					<i>Paramphistomum</i> spp.	-	-	18.4			
<i>Trichuris ovis</i>	25.4	-	-											
<i>T. axei</i>	19.4	-	-	<i>C. ovis</i>	11.9	-	-	<i>Fasciola hepatica</i>	4.5%	-	-	Debre Berhan	2500-300	Njau et al. (1990)
<i>H. contortus</i>	2.9	-	-	<i>Moniezia</i> spp.	7.5	-	-							
<i>Ostertagia</i> spp.	1.5	-	-											
Ascaris spp.	1.6	2.5	2.8											
Strongyle spp.	70.2	78.4	47					<i>Fasciola</i> spp.	0	0	0.4	Dembi Dolo	1500-2000	Fikru et al. (2006)
Dictyocaulus spp.	4.3	1.2	0.4											
Trichuris spp.	4.5	9.8	1.6											
-	-	-	-	<i>C.tenuicollis</i> , <i>C.ovis</i>	45.69,	72.38,	-							
				Hydatid cysts	8.43	8.57	-					Dessie	2,400	Abebe et al. (2015)
					9.02	1.90	-							
				<i>C.tenuicollis</i>	22.8	26.4	-					Dire Dawa	1204	Endale et al. (2013)
Shistosoma spp.	2.3	-	24.6									Durbete	1600	Yirsaw and Zewdu (2015)
Strongyle spp.	34.06	54	-					<i>F. hepatica</i>	8.52	6.99	-	Gondar	1500-2000	Shimelis et al. (2011)
Trichuris spp.	4.8	3	-	<i>Monezia</i> Spp.	3.93	0	-	<i>Paramphistomum</i> spp.	0.87	0	-			
-	-	-	-	Hydatid cysts	-	-	38.8					Gondar, Finote Selam, Injibara	1917-2560	Nigatu (2010)
<i>H.contortus</i> .	A*	A*	-	<i>M. expansa</i>	61	53								
<i>T.axei</i>	A*	A*	-											
<i>T.vitrinus</i>	A*	A*	-	<i>Av.centripunctata</i>	20	21		<i>P.microbothrium</i>	25	21				
<i>Trichostrongylus</i> spp.	A*	A*	-											
<i>Nematodirus spathir</i>	A*	A*	-	<i>St.globipunctata</i>	24	27		<i>F.hepatica</i>	26	3		Haremaya, Harar, Dire Dawa, Jijiga	2000	Menkir et al. (2007)
<i>Cooperia curticei</i>	A*	A*	-	<i>St.hepatica</i>	39	36		<i>F.gigantica</i>	20	10				
Strongyloides spp.	A*	A*	-											
<i>Bunostomum</i> spp	A*	A*	-	<i>Cysticercus ovis</i>)	26	22		<i>Dicrocoelium dendriticum</i>	7	2				
<i>Oes.columbianum</i>	A*	A*	-											
<i>Oesophagostomum</i>	A*	A*	-	<i>C.tenuicollis</i>	79	53								
<i>Chabertia ovina</i>	A*	A*	-											
Trichuris spp	A*	A*	-	<i>E. granulosus</i> (cysts)	68	65								
<i>Haemonchus</i> spp.	51.6	64.7												
<i>Trichostrongylus</i> spp.	38.7	41.2										Hawassa	1790	Abebe et al. (2010)
<i>Oesophagostomum</i>	22.6	23.5												
Trichuris spp.	27.6	12.5												
<i>Bunostomum</i> spp.	9.7	11.8												
-	-	-	-	Hydatid cysts	-	-	6.9	6.9	-	-		Hawassa	1790	Feyesa et al. (2010)
<i>Haemonchus</i> spp.	81.1	76.5	-									Hawassa	1790	Thomas et al. (2007)
<i>T.axei</i>	47.2	39.4	-											
Teladorsagia spp.	19.4	20.5	-											

Strongyels spp.	-	-	32.4												
Strongyloid spp.	-	-	3.3												
Trichuris spp.	-	-	1.9	M. expansa	-	-	5.2	Fasciola spp.	-	-	23.3	Jimma	1750	Hailu et al. (2011)	
Ascaris spp.	-	-	2.4					Paramphistomum	-	-	48.6				
Capillaria spp.	-	-	1.4												
Nematodirus spp.	-	-	0.9												
Strongyles spp.	27.58	-	3.6												
Trichuris spp.	2.95	-	0.74	Moniezia sp.	4.18	-	0.4	F. hepatica	1.47	-	4.8	Kelala,	1580	Tesfaye (1998)	
Strongyloides spp.	3.20	-	0.6												
Nematodirus spp.	0.49	-	5.3												
-	-	-	-	Hydatid cysts	-	-	4.25	-	-	-	-	Kombolcha	500 - 1840	Fufa et al. (2012)	
-	-	-	-	Cysticercus bovis	-	-	6.74	-	-	-	-	Mekelle	2000-2200	Getachew and Ashwani (2013)	
-	-	-	-	Hydatid cyst (E.granulosus)	-	-	8.1	-	-	-	-	Mekelle, Adigrat, Axum Humera, Maiche, Shire	1400 - 2500	Kebede et al. (2009)	
Haemonchus spp.	90.82	96.5		M.expansa	27.6	20.6									
T.axei	55.8	64.3													
T.colubriformis	87.15	90		C. tenuicolis	32.8	34.0	-	-	-	-	-	Metehara, Semera, Jigjiga	2300	Abebe and Esayas (2001)	
Bunostomum spp.	38.97	35.2	-												
Strongyloides spp.	38.02	43.6		Avitellina spp.	33.7	35.1	-	-	-	-	-				
Oesophagostomm.	74.8	70.8													
Trichuris spp.	51.75	48.18		Stilesia spp.	31.6	28.8									
-	-	-	-	-	-	-		Fasciola spp.	-	-	21.9	Nekemete	2,088	Alula et al.(2013)	
Haemonchus spp.	91.2	82.9	-	-	-	-	-	-	-	-	-	Ogaden	1200	Kumsa and Wossene (2006)	
Trichostrongylus spp.	37.7	40.2	-	-	-	-	-	-	-	-	-				
-	-	-	-	C. bovis	-	-	0.0	-	-	-	-	Wolaita Soddo	2500	Regassa et al. (2009)	
-	-	-	-	Hydatid cyst	-	-	9.9	-	-	-	-				
H.contortus	40.9	-	-	-	-	-	-	-	-	-	-	Wukro	1977	Lidya and Berihun (2015)	
- : No information available, A* = no specific prevalence is mentioned															

Species composition and seasonal dynamics

In Ethiopia, the presence of GI helminthes infections in domestic ruminants is reported by many authors (Thomas et al., 2007; Menkir et al., 2007; Abebe et al., 2010; Shimelis et al. 2011; Diriba et al., 2013; Addisu and Berihu, 2014 and Cheru et al. 2014). According to the current review, about twenty three (23) GI helminthes species belonging to Nematode, Cestode and Trematode categories have been reported from infected cattle, sheep and goats (Table 2). Whereas, Menkir et al. (2007), Abebe et al. (2010), Hailu et al. (2011) and Shimelis et al. (2011) who reported that the most prevalent genera of GI helminthes were *Haemonchus*, *Trichostrongylus*, *Oesophagostomum*, *Nematodirus*, *Cooperia*, *Toxocara* and *Bunostomum* from Nematodes; *Monezia* and *Cyticercus* from Cestodes and *Fasciola*, *Paramphistomum* and *Shistosoma* from Trematode classes (Table 3).

Table 3 - Summary of Species compositions of GI helminthes of domesticated ruminants in Ethiopia.

Major Class	Species	Host	Predilection site(s)
Nematode	<i>Haemonchus contortus</i>	Cattle, sheep, goats	Abomasums
	<i>Haemonchus placei</i>	Cattle	Abomasums
	<i>Ostertagia circumcincta</i>	Cattle, Sheep, Goats	Abomasums
	<i>Trichostrongylus axei</i>	Cattle, Sheep, Goats	Small intestine
	<i>Trichostrongylus colubriformis</i>	Cattle, Sheep, Goats	Small intestine
	<i>Cooperia curticei</i>	Cattle, Sheep, Goats	Small intestine
	<i>Strongyloides papillosus</i>	Cattle, Sheep, Goats	Small intestine
	<i>Ostertagia</i> Spp.	Cattle, Sheep, Goats	Abomasum, Large intestine
	<i>Chabertiaovina</i>	Cattle, Sheep, Goats	Large intestine
	<i>Oesophagostomum columbianum</i>	Sheep, Goat	Large intestine
	<i>Strongyloides papillosus</i>	Cattle, Sheep, Goats	Small intestine
	<i>Trichuris ovis</i>	Cattle, Sheep, Goats	Large intestine
	<i>Nematodirus fillicollis</i>	Cattle, Sheep, Goats	Small intestine
	<i>Nematodirus spathiger</i>	Sheep	Small intestine
	<i>Trichostrongylus vitrinus</i>	Sheep, Goat	Small intestine
<i>Toxocara vitulorum</i>	Cattle (calves)	Small intestine	
Cestode	<i>Monezia expansa</i>	Sheep, Goat	Small intestine
	<i>Cysticercustenuicollis</i> (<i>Taenia hydatigena</i>)	Sheep, Goat	Omentum Mesenteries Peritoneum, Liver
	<i>Ecchnococcus granulossus</i>	Sheep, Goat	Omentum Mesenteries Peritoneum, Liver
	<i>Cysticercus ovis</i> (<i>Taenia ovis</i>)	Sheep, Goat	Omentum Mesenteries Peritoneum, Liver
	<i>Avitellina centripunctata</i> (<i>Av.centripunctata</i>)	Sheep, Goat	Small intestine
	<i>Stilesia globipunctata</i> (<i>St.globipunctata</i>)	Sheep, Goat	Small intestine
	<i>Stilesia hepatica</i> (<i>St. hepatica</i>)	Sheep, Goat	Liver, biliary ducts
Trematode	<i>Fasciola hepatica</i>	Sheep, Goat Cattle	Liver
	<i>Fasciola gigantica</i>	Sheep, Goat Cattle	Liver
	<i>Paramphistomum cervi</i>	Sheep, Goat Cattle	Rumen
	<i>Paramphistomum microbothrium</i>	Sheep, Goat Cattle	Rumen
	<i>Schistosoma bovis</i>	Sheep, Goat Cattle	Liver, intestine, mesenteric lymph nodes and mesenteric veins

According to this systemic review, GI helminthiasis in domestic ruminants is severe and increasingly become an important focusing area of research in the country situation. A study conducted in western Oromia on GI parasites showed that Strangles were the most prevalent parasites encountered in the area (Moti et al., 2013). Similar study by Shimelis et al. (2011); Cheru et al. (2014), Nejib et al. (2016) and Diriba et al. (2013) reported that Strongyles were the most prevalent parasites encountered in North Gondar, Debre Zeit (East Shoa zone), Arbaminch (GamoGofa zone) and Asella respectively. As it has been reported by Abebe et al. (2015); Endale et al. (2013); Feyesa et al. (2010); Nigatu (2010); Kebede et al. (2009); Nigatu et al. (2009); Regassa et al. (2009); Jibat et al. (2008); Menkir et al. (2007); Bekele et al. (1988), metacestodes (larval cestodes) *Cysticercus ovis* (*Taenia ovis*), *Cysticercus tenuicollis* (*T. hydatigena*) and Hydatid cysts (*Echinococcus granulossus*) are the most prevalent species in Eastern Ethiopia. On the other hand, gastrointestinal infections as a result of adult cestodes such as *Avitellina centripunctata*, *Moniezia expansa* and *Stilesia globipunctata*, and bile duct infections with *Stilesia hepatica* were frequently reported in different parts of the country (Abebe and Esayas, 2001; Etana, 2002; Menkir et al., 2007; Bersissa et al., 2011). Nonetheless, a higher prevalence of strongyles infection was recorded in the midland and highland than the lowland, and in wet season than the dry season. The mean fecal egg count was found to be significantly higher in the midland area and in wet season (Nejib et al., 2016). However, according to a study by Abebe and Esayas (2001) in the arid and semiarid zones of Eastern Ethiopia revealed that during the dry seasons of the year, a greater prevalence rates of GI helminthes were recorded in sheep and goats, 95.6 % and 100 % respectively. Furthermore, according to an abattoir survey by Menkir et al. (2007) at 4 abattoirs located in the semi-arid zone of Eastern Ethiopia, the mean burdens of adult nematodes were generally moderate in both sheep and goats and showed patterns of seasonal abundance that corresponded with the bi-modal annual rainfall pattern, with

highest burdens around the middle of the rainy season. There were significant differences in the mean worm burdens and abundance of the different nematode species between the four geographic locations, with worm burdens in the Haramaya and Harar areas greater than those observed in the Dire Dawa and Jijiga locations (Abebe and Esayas, 2001; Menkir et al., 2007; Abebe et al., 2010; Berssisa et al., 2011; Endale et al., 2013). The seasonality of the GI helminthes distribution is associated with the relative humidity and rainfall (Debela, 2002; Menkir, 2007). A number of reports throughout the country indicated that there are remarkable changes in faecal egg counts and prevalence of helminthes infection as a result of seasonal variation and seasonal rainfall pattern (Fikru et al., 2006; Menkir et al., 2007; Takele et al., 2013). Furthermore, relatively higher GI Helminthes egg counts were found in mid altitude and highland zones than in lowland due to the influence of existing fluctuations in geographic and climatic conditions between each zone (Demelash et al., 2006; Takele et al., 2013).

Epidemiological factors

The epidemiology of the GI helminthiasis relies on factors such as the infection pressure in the environment and the susceptibility of the host, species and pathogen factor (Tilahun, 1995). From epidemiological point of view, the infective stages which eventually become available to the host depend on the independent and interactive influences of several factors in the macro- and micro-environment (Urquhart et al., 1994; Woldemariam, 2005; Regassa et al., 2006; Takele et al., 2013). Parasitic, host and environmental factors are the most frequently reported determinants for the epidemiology of helminthes (Etana, 2002; Fikru et al., 2006; Ahmed et al., 2007; Menkir, 2007; Shimelis et al., 2011; Yirsaw and Zewdu, 2015). As stated by Shimelis et al. (2011), the prevalence of helminthiasis at species level was about 46.07% and 55% in sheep and goats, respectively. Almost similar report was documented in and around Ambo town with the proportion of 47.8% and 53.3% in sheep and goats, correspondingly (Temesgen and Walanso, 2015). Among the collected articles, the highest post mortem examination result was reported in Eastern Ethiopia (100%) (Abebe and Esayas, 2001). Strong association between GI helminthes and poor body condition was coupled with heavy intensity of infection in the majority of infected animals (Abebe et al., 2010; Diriba et al. 2013; Alula et al., 2013; Temesgen and Walanso, 2015). According to Abebe et al. (2010) and Cheru et al. (2014), the burden of GI parasites and total EPG was significant in different body conditions.

Host factor

Sex, age, breed, nutrition, physiological status and presence or absence of inters- infections aggravate the severity of infection (Demelash et al., 2006; Menkir et al., 2007). Clinical parasitic gastroenteritis has been reported in young animals whilst infections in mature animals are generally subclinical in nature (Thomas et al., 2007). The lower occurrence in adults has been attributed to immunological maturity as the animals grow and the increase in acquired resistance due to repeated exposure (Biffa et al., 2004). While, some local breeds are known to be genetically resistant to GIhelminthes infections than others (Tibbo, 2006). In Ethiopia, the local sheep breeds (Washera, Farta, Afar, Menz, Horo); Goat breeds (Begait, Abewrgelie, Keffa,) and Cattle breeds (Boran , Fogera, Raya, Horro, Abigar, Shekko, Arssi) are relatively resistant to GI worms than exotic breeds (Frisch and O'Neill, 1998; Negussie et al., 2000; Tibbo, M. 2006; Menkir et al., 2007; Solomon et al., 2009; Kebede et al., 2012). In addition, Moti et al. (2013) reported that Physiological status of ruminants like level of host immunity to the parasites is subjected to the number of eggs produced by adult female helminthes. In the same way, the females are readily infected and existing worm burdens become more active and increase egg spassed in the feces and develop Larvae (L₃) on the pasture (Woldemariam, 2005).

Environmental factor

As far as Ethiopia is among the tropical African countries, the temperature is permanently favorable for larval development in the environment. The favorable environment for larvae development is ranged at temperature about 10–36 °C and humidity proportion of 85% (Debela, 2002). In the arid tropical climates of lowland areas of the country has an environment which ranges from extensive pasturelands and browse plants to intensive grazing areas (Nejib et al., 2016). This environment is ranged from harsh to favorable for growth and survival of free-living stages of the GI helminthes (Tilahun, 1995; Debela, 2002).

Pathogen factor

The epidemiology of GI helminthes is also strongly influenced by host-parasite biology after infection has been occurred (Abebe et al., 2010; Diriba et al., 2013). Hypobiosis has been undergone by GI helminthic the abomasal or intestinal mucosae of the host (Cheru et al., 2014). Whereas, Abebe and Esayas (2001) stated that the immune status of the host influenced the rates of hypobiosis and usually arrested during external environments are unfavorable for the development and survival of eggs and larvae. Such development cycle usually coincides with the onset of rainy seasons and favourable period for larval development and transmission (Kumsa and Wossenie, 2006; Feyisa et al., 2010; Hailu et al., 2011; Nejib et al., 2016).

Currently applied diagnostic techniques

The diagnosis of helminthes of ruminants is based on demonstrating the presence of eggs or larvae in fecal samples or parasites recovered from the digestive tracts or other viscera of the animals (Hailu et al., 2011; Addisu and Berihun, 2014). Although a great variety of methods and modifications have been described for diagnosis, standardized techniques such as egg or larval counts, worm counts and pasture larval counts did not exist. Therefore, most diagnostic laboratories

as well as teaching and research institutions were applied their own set of protocols and test procedures (Kassai, 1999). Common diagnostic procedures for helminthes infections in Africa in general and in Ethiopia in particular are simple flotation, sedimentation, modified McMaster and faecal culture methods (Hansen and Perry, 1994; Kassai, 1999; Waller, 1997). Some nematode genera such as strongyloides produce eggs that are identical in appearance which couldn't be identified easily by faecal examination alone. So to identify these faecal cultures are required (Hansen and Perry, 1994; Urquhart et al., 1996; Kassai, 1999; Van Wyket et al., 2004). However, *Nematodirus*, *Strongyloides* and *Trichuris* species have eggs that can be differentiated by their distinct morphological features. Post-mortem examinations and identification of adult worms and arrested larvae in animals are the definitive means of identifying the parasite. Similar to faecal egg counts; there are many procedures that are described for post-mortem examination for nematode parasites (Hansen and Perry, 1994; Urquhart et al., 1996; Kassai, 1999).

Chemotherapies and control options

Effective helminthic control is a major element in ensuring the sustainability of animal production. The main aim of control is therefore to ensure that the biotic potential of a parasite is restrained at a level compatible with the biological requirements of economic livestock production (Waller, 1997). Since eradication of gastrointestinal parasites is not practical, only integrated control methods can be envisaged. Some of the basic principles include grazing management, acquisition of natural or artificially induced immunity, biological control and the judicious use of anthelmintic (Hazelby et al., 1994). The main methods for control of helminthic parasites are prophylactic treatment with anthelmintics combined with grazing management (Van Wyk et al., 1999). Despite the accumulation of drugs in animal products and undesirable effects on non-target organisms in the environment, together with an increase in anthelmintic resistance, the use of anthelmintics still remains the corner-stone of helminth control (Waller, 1997; Van Wyk et al., 1999; Bersissa and Girma, 2009). Since animals are often infected with a wide range of helminths, the need for broad-spectrum compounds against trematodes, cestodes and nematodes, and their larval stages is obvious (Hazelby et al., 1994). The epidemiological information on GI helminths parasites of domestic ruminants gathered in Ethiopia can be used to design appropriate control measures. In principle, control should aim at the reduction of transmission rates. Several control methods, which include cultural husbandry, chemical, biological, ethno-veterinary medicine and immunological control, have been proposed (Nejib et al., 2016).

Husbandry pattern, control and prevention

A thorough husbandry practices such as controlling stocking rates, rotational grazing, and providing hygienic grazing can be considered as an alternative husbandry control technique (Diriba and Birhanu, 2013; Abebe et al., 2015; Temesegegn and Walanso, 2015). The best way to prevent GI helminthes is to keep animals away from potentially dangerous environment. An absolute separation of stock from intermediate host zone is only practical in intensive farming husbandry systems (Woldemariam, 2005) which the country has a limited effort to do so. However, in communal grazing condition which is very common and traditional in Ethiopia, animals are communally grazed and therefore; practices such as rotational grazing and provision of clean pastures would not be feasible (Menkir et al., 2007; Cheru et al., 2014).

Chemotherapeutic Interventions

Nowadays, the control of GI parasites of livestock in Ethiopia is mainly based on the use of anthelmintics. The most commonly used generic broad spectrum anthelmintics that are available in Ethiopia are the Benzimidazoles, Imidazothiazoles and Macrocytic lactones which consist albendazole, levamisole and ivermectin, respectively (Woldemariam, 2005; Kumsa, and Wossene, 2006; Menkir et al., 2007). It is readily available wherever in the country since it has been imported massively by the government and non-government institutions; and used by every individual including farmer. However, the use of anthelmintics at regular intervals for a long period of time and treating by mass whenever an animal manifest clinical syndrome has become the major issue for the development of multiple resistances; for instance against benzimidazoles (Woldemariam, 2005). Targeted or selective application of anthelmintic treatment might be an important tool to keep susceptible GI nematode strains in livestock and to delay this case. An alternative approach for selective anthelmintic treatment was studied using experimental small ruminants for the management of haemonchosis by using the FAMACHA® method in the Mid-Rift Valley of Ethiopia by Woldemariam (2005) and Menkir (2007). Such system can be used by the farmers themselves by checking their animals for signs of anaemia (VanWyket al., 2004). Unfortunately, it was not practiced very well in Ethiopia where mixed parasitic infection, and where traditional feeding practice is followed (VanWyk et al., 2004; Kumsa and Wossene, 2006). A cost-effective preventive control programs for helminth infection in ruminants is based on sound epidemiological knowledge of the time relationship between contamination of pastures and the seasonal availability of infective larvae in a given geographic area. Epidemiological knowledge, its application in grazing management whenever feasible and access to anthelmintics of high efficacy are key factors for the success of controlling helminth infections in domestic ruminants (Aynalem et al., 2009; VanWyk and Mayhew, 2013). Projecting models derived from more complete information on the ecology of GI helminth infection and anthelmintic resistance, climate and local management factors provide a basis for improved control schemes based on chemotherapy, management and immunization within similar climate (VanWyk and Mayhew, 2003; Woldemariam, 2005; Menkir et al., 2007).

Ethno-veterinary practices

As a result of the gradually increasing anthelmintic resistance, residual effect on animal products, environmental pollution, scarcity and high cost of such drug especially to poor farmers have enforced to reconsider other alternative helminthic control technique in the country (Bersissa and Girma, 2009). Of these, Ethno-veterinary medicine has become a substantial and most expanding interest of options for Ethiopian farmers. Although such a kind of conventional veterinary medical system was yet very poor in the country, a very limited effort have been done to encourage the widely used ethno-veterinary plants in the country (Lulekal et al., 2008; Fullas, 2010; Asfaw and Fentahun, 2020). In order to do so, several Ethno-veterinary surveys were conducted so far in the country which indicated as if several traditional healers use medicinal plants for de-worming livestock (Jemal et al., 2011). To mention few examples, Herbal preparations from fresh leaves of *Dodonea viscosa*, *Albizia gummifera* and *Vernonia amygdalina* against mixed natural infections in sheep was evaluated by Biffa et al. (2004) to show the anthelmintic activities. In addition, an In-vitro antihelmintics activity study from *Rhus glutinosa*, *Syzygium guineense*, *Albizia gummifera*, *Croton macrostachyus*, *Ekebergia capensis*, *Acacia nilotica* and *Terminalia schimperiana* against *Haemonchis contortus* have been reported by Eguale et al. (2006) and Jemal et al. (2011). Furthermore, anthelmintic activity of plants such as *Allium sativum*, *Zingiber officinale*, *Cucurbita mexicana*, *Ficus religiosa*, *Artemisia brevifolia*, *Calotropis procera*, *Nicotiana tabacum*, *Butea monosperma*, *Coriandrum sativum*, *Ocimum*, *Thymus schimperi* and *Echinops kebericho* have been reported by Abera (2003), Biffa et al. (2004) and Giday et al. (2007).

CONCLUSION

This review work assessed the GI helminthes in Ethiopia and provides a clue on perspectives and constraints encountered in researches which were done on GI helminths in ruminants. Twenty three (23) GI helminthes species that belong to all the three major classes of helminths have been found to occur in domestic ruminants in Ethiopia. In addition, nematodes are the most commonly encountered GI helminths while Cestodes are the least. In most reports, a higher rate was recorded in small ruminants. The most prevalent genera of GI helminths reported in order of prevalence are *Haemonchus*, *Trichostrongylus*, *Oesophagostomum*, *Nematodirus*, *Cooperia*, *Toxocara* and *Bunostomum* from Nematodes; *Monezia* and *Cycticercus* from Cestodes. Whereas, *Fasciola*, *Paramphistomum* and *Shistosoma* are found from Trematode category. Both the abattoir and coprological studies have indicated that infection by GI helminthes in ruminants is highly prevalent and widespread in all agro-ecologies and livestock production systems in Ethiopia. Fecal diagnostic techniques such as simple flotation, sedimentation, modified McMaster have been used routinely in Ethiopia. It has been also shown that prevalence of GI helminthes parasites was related to the agro-climatic conditions such as quantity and quality of pasture, temperature, humidity and grazing behavior of the host and the susceptibility of any intestinal helminthic parasites were also influenced by age, breed, species, health status, physiological factors and previous exposure to parasites.

Due to the lack of effective helminthes control strategies in Ethiopia, antihelmintics are exclusively used. Though Ethiopia has a huge amount of small and large ruminants population, the country is facing a direct and indirect economic lose as a result of GI helminthes infection. Hence, immediate remedies shall be taken into action on control and prevention methods against such anthelmintic resistant GI parasites. In order to this, it is advantageous to collect and looking over the previous researches done so far to reconsider their gaps for the future short and long term actions on prevention and control strategies. So, all-inclusive and well organized documentation about GI helminthes of ruminants in the country is essential to support researchers and policy makers to develop such remedies. Finally, applicable field diagnostic technique should be introduced as far as mixed parasitic infection and traditional feeding practice is common in the country which can aggravate GI helminthes infection.

DECLARATIONS

Consent to publish

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by the University of Gondar.

Availability of data and materials

Data will be made available up on request of the primary author

Acknowledgment

The authors' heartfelt thanks the University of Gondar, Research and Community Service V/President for the financially supporting in the study.

REFERENCES

- Abebe R, Mebrahtu G, Solomon M, Fufa A, and Alemayehu R (2010). Gastrointestinal nematode infections in small ruminants under the traditional husbandry system during the dry season in southern Ethiopia. *Tropical Animal Health and Production*, 42:1111–1117. Doi: <https://doi.org/10.1007/s11250-010-9532-3>
- Abebe TG, Belay M, Shahid N and Assefa A (2015). Major metacestodes in small ruminants slaughtered at Dessie municipal abattoir, Eastern Ethiopia: Prevalence, cyst viability, organ distribution and economic implications. *Comparative Clinical Pathology*, 24: 659–668. Doi: <https://doi.org/10.1007/s00580-014-1964-0>.
- Abebe W, and Esayas G (2001). Survey of Ovine and Caprine Gastro-intestinal helminthosis in eastern part of Ethiopia during the dry season of the year. *Revue de Médecine Vétérinaire*, 152 (5), 379-384. Link: https://www.revmedvet.com/2001/RMV152_379_384.pdf
- Abera B (2003). Medicinal plants used in traditional medicine in Jimma Zone, Southwest Ethiopia. *Ethiopian Journal of Health Sciences*, 13: 85–94. Link: <https://www.ajol.info/index.php/ejhs/article/view/146241>
- Addisu B, and Berihu H (2014). Study on Prevalence of Gastrointestinal Nematodes and Coccidian Parasites Affecting Cattle in West Arsi zone, Oromia Regional State, Ethiopia. *Journal of Veterinary Science and Technology*, 5(5): 1000207. Link
- Ahmed EF, Markvichitr K, Tumwasorn S, Koonawootrittriron S, Choothesa A, Jittapalpong S (2007). Prevalence of *Fasciola* spp. Infection of sheep in the Middle Awash River Basin, Ethiopia. *Southeast Asian Journal of Tropical Medicine and Public Health*, Vol 38 (suppl 1): 51-57. Link: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.1067.4234&rep=rep1&type=pdf>
- Alula P, Addisu K, and Amanuel W (2013). Prevalence and economic significance of Bovine Fasciolosis in Nekemte Municipal abattoir. *Journal of Veterinary Medicine and Animal Health*, 5(8): 202-205. Link: https://academicjournals.org/article/article1379686620_Petros%20et%20al.pdf
- Asfaw M and Fentahun T (2020). Treatment trials of epizootic lymphangitis with local medicinal plants: a review. *Online Journal of Animal and Feed Research*, 10(4): 93-101. Link
- Ayalew S, and Endalkachew N (2013). Prevalence and Risk Factors of Bovine and Ovine Fasciolosis, and evaluation of direct sedimentation sensitivity method at Bahir Dar Municipal Abattoir, Northern Ethiopia. *Ethiopian Veterinary Journal*, 17(2):1-17. Doi: <https://doi.org/10.4314/evj.v17i2.1>
- Aynalem H, Abebaw G, Taye T, and Markos T (2009). Epidemiology of nematode parasites of sheep around Jimma, southwestern Ethiopia. *Tropical Animal Health and Production*, 42: 825–831. Doi: <https://doi.org/10.1007/s11250-009-9492-7>
- Bekele T, Mukasa-Mugerwa E and Kasali OB (1988). The prevalence of cysticercoids and hydatidosis in Ethiopian sheep. *Veterinary Parasitology*, 28: 267-270. Doi: [https://doi.org/10.1016/0304-4017\(88\)90114-8](https://doi.org/10.1016/0304-4017(88)90114-8)
- Bekele T, Woldeab T, Lahlou-Kassi A, and Sherngton J (1992). Factors affecting morbidity and mortality on farm and on station in Ethiopia highland sheep. *Acta Tropica*, 52: 99-109. Doi: [https://doi.org/10.1016/0001-706X\(92\)90025-S](https://doi.org/10.1016/0001-706X(92)90025-S)
- Bersissa K and Girma A (2009). Multiple anthelmintic resistance on a goat farm in Hawassa (southern Ethiopia). *Tropical Animal Health and Production*, 41: 655–662. Doi: <https://doi.org/10.1007/s11250-008-9237-z>
- Bersissa K, Tigist T, Teshale S, Reta D, and Bederu H (2011). Helminthes sheep and goats in central Oromia (Ethiopia). During the dry season. *Journal of Animal and Veterinary Advances*, 10(14): 1845-1849. Link
- Beyene T and Hiko A (2019). Zoonotic metacestodes and associated financial loss from cattle slaughtered at Yabello municipal abattoir, Borana-Oromia, Ethiopia. *Parasite Epidemiology and Control*, 5: e00096. Doi: <https://doi.org/10.1016/j.parepi.2019.e00096>
- Biffa D, Nurfeta A, and Jobre Y (2004). Evaluation of anthelmintic activities of crude leaf extracts of three indigenous herbal plants against ovine gastrointestinal nematodes. *Ethiopian Veterinary Journal*, 8: 57-58.
- Cheru T, Birhanu A, Diriba L, and Eyob E (2014). Prevalence of gastrointestinal parasitism of cattle in East Showa Zone, Oromia Regional State, Central Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 6(2): 54-62. Link: <https://academicjournals.org/journal/JVMAH/article-full-text-pdf/D0DC71C42793>
- CSA (2013). Agricultural Sample Survey, 2012/13, Volume II: Report on Livestock and livestock characteristics (Private peasant holdings). Statistical Bulletin 570. Central Statistical Agency (CSA), Federal Democratic Republic of Ethiopia, Addis Ababa.
- Debela E (2002). Epidemiology of gastrointestinal helminthiasis of Riff valley goats under traditional husbandry system in Adami Tulu district. Ethiopia. *Ethiopian Journal of Sciences*. 25, 35-44. Link: <https://www.ajol.info/index.php/sinet/article/view/18072>
- Demelash B, Yilma J, and Hassen C (2006). Ovine helminthosis, a major health constraint to productivity of sheep in Ethiopia. *Animal Health Research Reviews*, 7(1/2), 107–118. Doi: <https://doi.org/10.1017/s1466252307001132>
- Diriba L, and Birhanu A (2013). Prevalence of ovine gastrointestinal nematodes in and around Asella, South Eastern Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 5(8): 222-228. Link
- Egualé T, Tilahun G, Gidey M, and Mekonnen Y (2006). In-Vitro anthelmintic activities of four ethiopian medicinal plants against *Haemonchus contortus*. *Pharmacologyonline*, 3:153-165. Link | [Google Scholar](#)
- Elsa L, Sofia V, Elvira S, Maria M, Mendes G, and André M (2012). Factors Influencing Livestock Productivity. *Environmental Stress and Amelioration in Livestock Production*, Springer-Verlag Berlin Heidelberg. Link
- Endale M, Shihun S, Jemere B, and Desie S (2013). Sheep and goats *Cysticercus tenuicollis* prevalence and associated risk factors. *African Journal of Agricultural Research*, 8(24): 3121-3125. doi: <https://doi.org/10.5897/AJAR2012.7361>
- Etana D (2002). Epidemiology of Gastrointestinal helminthes of Rift valley goats under traditional husbandry systems in Adami Tulu District, Ethiopia. *Ethiopian Journal of Science*, 25(1): 35-44. Link: <https://www.ajol.info/index.php/sinet/article/view/18072>
- Feyesa R, Alemante M, and Jemere B (2010). Study on the prevalence of cystic hydatidosis and its economic significance in cattle slaughtered at Hawassa Municipal abattoir, Ethiopia. *Tropical Animal Health and Production*, 42: 977–984. Doi: <https://doi.org/10.1007/s11250-009-9517-2>
- Fikru R, Yosef K, Teshale S, Reta D, and Yosef K (2006). Epidemiology of Gastrointestinal Parasites of Ruminants in Western Oromia, Ethiopia. *International Journal Applied Research and Veterinary Medicine*, 4:51-57. Link
- Frisch JE, and O'Neill CJ (1998). Comparative evaluation of beef cattle breeds of African, European and Indian origins. 1. Live weights and heterosis at birth, weaning and 18 months. *Animal Science*, 67: 27-38. Doi: <https://doi.org/10.1017/S135772980009760>
- Fufa A, Sisay F, Bekele M, and Alemayehu R (2012). Prevalence of bovine hydatidosis in Kombolcha ELFORA abattoir, North Eastern Ethiopia. *Open Journal of Animal Sciences*, 4:281-286. <http://dx.doi.org/10.4236/ojas.2012.24038>.
- Fullas F (2010). Ethiopian medicinal plants in veterinary healthcare. A mini-review. *Ethiopian eJournal for Research and Innovation Foresight*, 2: 48–58. Link

- Getachew A and Ashwani K (2013). Cysticercosis in cattle and its public health implications in Mekelle City and surrounding areas, Ethiopia. *Ethiopian Veterinary Journal*, 17: 31-40. Link: <https://www.ajol.info/index.php/evj/article/download/99352/88643>
- Getawa A, Beyene D, Ayana D, Megersa B, and Abunna F (2010). Hydatidosis: Prevalence and its economic importance in ruminants slaughtered at Adama municipal abattoir, Central Oromia, Ethiopia. *Acta Tropica*, 221–225. Doi: <https://doi.org/10.1016/j.actatropica.2009.10.019>
- Giday M, Teklehaymanot T, Anmut A, and Mekonnen Y (2007). Medicinal plants of the Shinasha, Agew Awi and Amhara peoples in northwest Ethiopia. *Journal of Ethnopharmacology*, 110: 516–525. Doi: <https://doi.org/10.1016/j.jep.2006.10.011>
- Hailu D, Cherenet A, Moti Y, and Tadele T (2011). Gastrointestinal Helminth Infections in Small-Scale Dairy Cattle Farms of Jimma Town, Ethiopia. *Ethiopian Journal of Applied Sciences and Technology* 2(1): 821. [Link](#)
- Hansen J, and Perry B (1994). The epidemiology diagnosis and control of helminth parasites of ruminants. A Handbook. International Laboratory for Research on Animal Diseases, Nairobi, Kenya. Link: <https://cgspace.cgiar.org/handle/10568/49809>
- Hazelby, CA, Probert, AJ, and Rowlands DAP (1994). Anthelmintic resistance in nematodes causing parasitic gastroenteritis of sheep in the UK. *Journal of Veterinary Pharmacology and Therapeutics*. 17(4): 245–252. Doi: <https://doi.org/10.1111/j.1365-2885.1994.tb00240.x>
- Jemal H, Kaba U, Fayissa R, Awol J, Sultan A, and Nezif H (2011). Antihelmintic Effect of the Essential Oil Extracts of Selected Medicinal Plants against *Haemonchus contortus*. *International Journal of Agricultural Research*, 6: 290-298. Doi: <http://dx.doi.org/10.3923/ijar.2011.290.298>
- Jibat T, Ejeta G, Asfaw Y, and Wudie A (2008). Causes of abattoir condemnation in apparently healthy slaughtered sheep and goats at HELMEX abattoir, Debre Zeit, Ethiopia. *Revue de Médecine Vétérinaire*, 159(5): 305-311. Link: https://www.revmedvet.com/2008/RMV159_305_311.pdf
- Kassai T (1999). *Veterinary Helminthology*. Reed Educational and professional publishers, Oxford UK, 260.
- Kebede T, Haile A, Dadi H (2012). Smallholder goat breeding and flock management practices in the central rift valley of Ethiopia. *Tropical Animal Health and Production*, 44: 999–1006. Doi: <https://doi.org/10.1007/s11250-011-0033-9>
- Kebede W, Hagos A, Girma Z, Lobago F (2009). Echinococcosis/hydatidosis: its prevalence, economic and public health significance in Tigray region, North Ethiopia. *Tropical Animal Health and Production*, 41: 865–871. Doi: <https://doi.org/10.1007/s11250-008-9264-9>
- Khan MK, Sajid MS, Khan MN, Iqbal Z, Iqbal MU (2009). Bovine Fasciolosis: Prevalence, effects of treatment on productivity and cost benefit analysis in five districts of Punjab, Pakistan. *Research in Veterinary Science*, 87: 70–75. Doi: <https://doi.org/10.1016/j.rvsc.2008.12.013>
- Kumsa B, and Wossene A (2006). Abomasal nematodes of small ruminants of Ogaden region, eastern Ethiopia: Prevalence, worm burden and species composition. *Revue de Médecine Vétérinaire*, 157(12): 27-32. [Link](#)
- Lebbie SH, Rey B, and Irungu EK (1994). Small ruminant research and development in Africa Proceedings of the second biennial conference of the African small ruminant research network. ILCA, pp.1-5. [Link](#)
- Lemma B, Gebreab F, and Tedla S (1985). Studies on fascioliasis in four selected sites in Ethiopia. *Veterinary Parasitology*, 18: 29-37. DOI: [https://doi.org/10.1016/0304-4017\(85\)90005-6](https://doi.org/10.1016/0304-4017(85)90005-6)
- Lidya G, and Berihun AT (2015). Prevalence of Ovine Haemonchosis in Wukro, Ethiopia. *Hindawi Publishing Corporation Journal of Parasitology Research*, Article ID: 635703. Doi: <http://dx.doi.org/10.1155/2015/635703>.
- Lulekal E, Kelbessa E, Bekele T, and Yineger H (2008). An ethno-botanical study of medicinal plants in Mana Angetu District, southeastern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 4:10. Doi: <https://doi.org/10.1186/1746-4269-4-10>
- Menkir MS, Arvid U, and Peter Waller J (2007). Prevalence and seasonal incidence of nematode parasites and fluke infections of sheep and goats in eastern Ethiopia. *Tropical Animal Health and Production*, 39: 521–531. Doi: <https://doi.org/10.1007/s11250-007-9035-z>
- Menkir S (2007). *Helminth Parasites of Sheep and Goats in Eastern Ethiopia: Epidemiology, and Anthelmintic Resistance and its Management*. Doctoral thesis, Faculty of Veterinary Medicine and Animal Science, Swedish University of Agricultural Sciences, Uppsala, Sweden. Link: <https://core.ac.uk/download/pdf/11695158.pdf>
- Moti Y, Derara B, Dalasa D, Senbeta T, and Hailu D (2013). Bovine Trypanosomosis and gastrointestinal helminthosis in settlement villages of Bedele district, South-western Ethiopia. *Ethiopian Veterinary Journal*, 17(1):41-54. Doi: <https://doi.org/10.4314/evj.v17i1.4>
- Negussie E, Rottman OJ, Pirchner F, Rege JE (2000). Allometric growth coefficients and partitioning of fat depots in indigenous Ethiopian Menz and Horro sheep breeds. In *The Opportunities and Challenges of Enhancing Goat Production in East Africa*. Workshop Proceedings. Langston University, OK (USA). E (Kika) dela Garza Inst. for Goat Research. Langston, OK (USA) 2000 (pp. 151-163). [Google Scholar](#)
- Nejib M, Messele T, Amenu A, and Desie S (2016). Epizootological study of small ruminant gastrointestinal strongyles in Gamo-Gofa Zone, Southern Ethiopia. *Journal of Parasitic Disease*, 40(2): 469–474. DOI: <https://dx.doi.org/10.1007%2Fs12639-014-0528-1>
- Nigatu K, Abebe M, and Getachew T (2009). Hydatidosis of slaughtered animals in Bahir Dar Abattoir, Northwestern Ethiopia. *Tropical Animal Health and Production*, 41:43–50. Doi: <https://doi.org/10.1007/s11250-008-9152-3>
- Njau BC, Scholtens RG and Kasali O (1990). Parasites of sheep at the International Livestock Centre for Africa Debre Berhan Station, Ethiopia. *Preventive Veterinary Medicine*, 9: 267-277. Doi: [https://doi.org/10.1016/0167-5877\(90\)90072-P](https://doi.org/10.1016/0167-5877(90)90072-P)
- Regassa A, Abunna F, Mulugeta A, and Megersa B (2009). Major metacestodes in cattle slaughtered at Wolaita Soddo Municipal abattoir, Southern Ethiopia: Prevalence, cyst viability, organ distribution and socioeconomic implications. *Tropical Animal Health and Production*, 41: 1495–1502. Doi: <https://doi.org/10.1007/s11250-009-9338-3>
- Regassa F, Sori T, Dhuguma R, and Kiros Y (2006). Epidemiology of gastrointestinal parasites of ruminants in western Oromia, Ethiopia. *International Journal Applied Research of Veterinary Medicine*, 4(1): 51-57. Doi: <https://www.jarvm.com/articles/Vol4Iss1/Vol4Iss1RegassaV4N1pp51-57.pdf>
- Shimelis D, Asmare A, and Wudu T (2011). Epidemiology of gastrointestinal helminthiasis of small ruminants in selected sites of North Gondar zone, Northwest Ethiopia. *Ethiopian Veterinary Journal* 15 (2): 57-68. Doi: <https://doi.org/10.4314/evj.v15i2.67694>
- Solomon G, Komen V, Arendonk JAM (2009). Optimal village breeding schemes under smallholder sheep farming systems. *Livestock Science*, 124: 82–88. Doi: <https://doi.org/10.1016/j.livsci.2009.01.001>
- Takele S, Yacob H, and Getachew T (2013). Epidemiology of gastrointestinal nematodes of Horro sheep in Western Oromiya, Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 5(10): 296-304. Doi: <https://doi.org/10.5897/JVMAH2013.0234>

- Tembely S, and Hansen JW (1996). Helminth Diseases of Small Ruminants in the Tropics: A Review of Epidemiology and Control Strategies, Bogor, Indonesia, 1996: 123-127. [Link](#)
- Temesgen A, and Walanso I (2015). Major gastrointestinal helminth parasites of grazing small ruminants in and around Ambo town of Central Oromia, Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 7: 64-70 . Doi: <https://doi.org/10.5897/JVMAH2014.%200327>
- Tesfaye H (1998). Ovine and bovine helminthiasis in Kelala (South Wollo). *Ethiopian Vet Association Proceedings of the 12th Conference*. Addis Ababa, Ethiopia, pp. 30-34.
- Thomas N, Teshale S, and Kumsa B (2007). Abomasal nematodes of sheep and goats slaughtered in Awassa (Ethiopia): Species composition, prevalence and vulvar morphology. *Helminthologia*, 44: 70–75. Doi: <https://doi.org/10.2478/s11687-007-0006-8>
- Tibbo M (2006). Productivity and health of indigenous sheep breeds and crossbreds in the central Ethiopian highlands. Doctoral dissertation. ISSN 1652-6880, ISBN 91-576-7100-1. Link: <https://core.ac.uk/download/pdf/11695029.pdf>
- Tilahun G (1995). Epidemiology of helminth parasites of small ruminants in midlowland Ethiopia. In: Gray GD, Uilenberg G, Jacobsen LL (Eds.), *Parasitology Research in Africa. Proceedings of an IFS (International Foundation for Science) workshop*, Bobo Dioulasso, Burkina Faso. pp. 255-269. [Google Scholar](#)
- Tilahun H, and Schmidt E (2012). Spatial analysis of livestock production patterns in Ethiopia. ESSP II Working Paper 44. International Food Policy Research Institute/Ethiopia Strategy Support Program II, Addis Ababa, Ethiopia. Link: <https://www.ifpri.org/publication/spatial-analysis-livestock-production-patterns-ethiopia>
- Urquhart GM, Armour J, Duncan JL, Dunn AM, Jennings FW (1996). *Veterinary Parasitology* 2nd Edn, Oxford, Longman Scientific and Technical Press, UK. 100-109.
- Van Wyk JA, and Mayhew E (2013). Morphological identification of parasitic nematode infective larvae of small ruminants and cattle: A practical lab guide. *Onderstepoort Journal of Veterinary Research*, 80: E1-E14. [Link](#)
- Van Wyk JA, Cabaret J, and Michael LM (2004). Morphological identification of nematodes of small ruminants and cattle simplified. *Veterinary Parasitology*, 119: 277–306. Doi: <https://doi.org/10.1016/j.vetpar.2003.11.012>
- Van Wyk JA, Malan FS and Randles JL (1999). How long before resistance makes it impossible to control some field strains of *Haemonchus contortus* in South Africa with any of the modern anthelmintics. *Veterinary Parasitology*, 70: 111-122. Doi: [https://doi.org/10.1016/S0304-4017\(96\)01147-8](https://doi.org/10.1016/S0304-4017(96)01147-8)
- Waller PJ (1997). Sustainable helminth control of ruminants in developing countries. *Veterinary Parasitology*, 71: 195-207. Doi: [https://doi.org/10.1016/s0304-4017\(97\)00032-0](https://doi.org/10.1016/s0304-4017(97)00032-0)
- Winter J, Rehbein S, and Joachim A (2018). Transmission of helminthes between species of ruminants in Austria appears more likely to occur than generally assumed. *Frontiers in Veterinary science*, 5: 30. Doi: <https://doi.org/10.3389/fvets.2018.00030>
- Woldemariam DL (2005). Nematode prevalence, helminths management practices and anthelmintic resistance in small ruminants in the mid-rift valley of Ethiopia. Thesis, Faculty of Veterinary Science, and University of Pretoria, South Africa. [Google Scholar](#)
- Yirsaw K, and Zewdu S (2015). Bovine and ovine schistosomiasis: prevalence and associated risk factors in South Achefer. The 25th Staff and students annual research conference, June 19-20, 2015. Gondar, Ethiopia. [Google Scholar](#)
- Yitayal G, Mebrehatu G, and Abebe T (2015). Prevalence of Bovine Fasciolosis in and around Bahir Dar, Northwest Ethiopia. *Journal of Parasitology and Vector Biology*, 7(4): 74-79. Link: https://academicjournals.org/article/article1429098988_Gebrie.pdf
- Zahid IA, Latif M, and Baloch KB (2005). Incidence of endoparasites in exotic cattle calves. *Pakistan Veterinary Journal*, 25 (1): 47-48. [Link](#)